

Gauge theory, line operators and dualities

Filippo Passerini
CERN

GeNeZiSS String Theory Meeting
École Polytechnique Fédérale de Lausanne

November 30, 2012

My Interests:

- formal aspects of supersymmetry
- non-local operators in gauge theory
- interconnections between gauge theory, string theory (or M-theory), 2d CFT and matrix models

AdS/CFT: The strong coupling regime of a gauge theory is encoded by the dynamics of weakly coupled strings in a certain background.

$SU(N)$ $\mathcal{N}=4$ SYM: the dual string theory is IIB on $AdS_5 \times S^5$ with tension $T = \frac{\sqrt{\lambda}}{2\pi}$, where $\lambda = Ng^2$. The conformal 4d gauge theory lives on the boundary of AdS_5

AdS/CFT: The strong coupling regime of a gauge theory is encoded by the dynamics of weakly coupled strings in a certain background.

$SU(N)$ $\mathcal{N}=4$ SYM: the dual string theory is IIB on $AdS_5 \times S^5$ with tension $T = \frac{\sqrt{\lambda}}{2\pi}$, where $\lambda = Ng^2$. The conformal 4d gauge theory lives on the boundary of AdS_5

SUSY Wilson Loops

$$W_R(C) = \frac{1}{N} \text{tr}_R \mathcal{P} \exp \left[\int_C ds (iA_\mu(x)\dot{x}^\mu + n_I \Phi_I(x)|\dot{x}|) \right]$$

In the string theory dual, the Wilson loop is associated to an open string that ends on the boundary of AdS_5 , on the loop where the operator is supported

[Maldacena] [Rey, Yee]

$$\langle W(C) \rangle = \int_{\partial X=C} DX e^{-T S[X]}$$

AdS/CFT: The strong coupling regime of a gauge theory is encoded by the dynamics of weakly coupled strings in a certain background.

$SU(N)$ $\mathcal{N}=4$ SYM: the dual string theory is IIB on $AdS_5 \times S^5$ with tension $T = \frac{\sqrt{\lambda}}{2\pi}$, where $\lambda = Ng^2$. The conformal 4d gauge theory lives on the boundary of AdS_5

SUSY Wilson Loops

$$W_R(C) = \frac{1}{N} \text{tr}_R \mathcal{P} \exp \left[\int_C ds (iA_\mu(x)\dot{x}^\mu + n_I \Phi_I(x)|\dot{x}|) \right]$$

In the string theory dual, the Wilson loop is associated to an open string that ends on the boundary of AdS_5 , on the loop where the operator is supported

[Maldacena] [Rey, Yee]

$$\langle W(C) \rangle = \int_{\partial X=C} DX e^{-T S[X]}$$

Strong coupling $\lambda \gg 1$ and $N \rightarrow \infty$, semiclassical string

[Drukker, Gross]


$$\langle W(C_{\text{circle}}) \rangle_{\mathcal{N}=4} \simeq KT^{-3/2} e^{2\pi T}$$

In agreement with an exact gauge theory computation using localization

[Pestun]

Next Step: reduce the symmetry \Rightarrow increase complexity of the theory

$\mathcal{N} = 2$ $SU(N)$ SCYM:




Interesting features: matter fields, non-trivial instanton dynamics, ...

The string dual is not known for this $\mathcal{N} = 2$ theory.

Next Step: reduce the symmetry \Rightarrow increase complexity of the theory

$\mathcal{N} = 2$ $SU(N)$ SCYM:




Interesting features: matter fields, non-trivial instanton dynamics, ...

The string dual is not known for this $\mathcal{N} = 2$ theory.

The vev of certain observables can be computed exactly using localization [Pestun]
 \rightarrow use these results to get information about the string dual of $\mathcal{N} = 2$ SCYM !

Next Step: reduce the symmetry \Rightarrow increase complexity of the theory

$\mathcal{N} = 2$ $SU(N)$ SCYM:



Interesting features: matter fields, non-trivial instanton dynamics, ...

The string dual is not known for this $\mathcal{N} = 2$ theory.

The vev of certain observables can be computed exactly using localization [Pestun]
 \rightarrow use these results to get information about the string dual of $\mathcal{N} = 2$ SCYM !

Pestun Localization: reduce the field theory path integral to a matrix integral, for any value of the gauge coupling. VEV of certain non-local operators can be computed using a matrix model.

$$Z = \int D\Psi e^{-S[\Psi]} = \int DM e^{-S[M]} Z_{1\text{-loop}}(M) Z_{\text{inst}}(M)$$

- M is a constant value of an adjoint scalar, i.e. a matrix

$\frac{1}{2}$ BPS Wilson loop can be computed as an observable of the matrix model

$$\langle W_R(\text{Circle}) \rangle = \left\langle \frac{1}{N} \text{tr}_R e^{2\pi M} \right\rangle_{\text{Matrix Model}}$$

$\frac{1}{2}$ BPS Wilson loop can be computed as an observable of the matrix model

$$\langle W_R(\text{Circle}) \rangle = \left\langle \frac{1}{N} \text{tr}_R e^{2\pi M} \right\rangle_{\text{Matrix Model}}$$

Strong coupling $\lambda \gg 1$, Large N limit: $N \rightarrow \infty$ with $\lambda = Ng^2$ fixed [Passerini, Zarembo]

$$\langle W(C_{\text{circle}}) \rangle_{\lambda \gg 1} = \text{const} \frac{\lambda^3}{(\ln \lambda)^{3/2}}$$

$\frac{1}{2}$ BPS Wilson loop can be computed as an observable of the matrix model

$$\langle W_R(\text{Circle}) \rangle = \left\langle \frac{1}{N} \text{tr}_R e^{2\pi M} \right\rangle_{\text{Matrix Model}}$$

Strong coupling $\lambda \gg 1$, Large N limit: $N \rightarrow \infty$ with $\lambda = Ng^2$ fixed [Passerini, Zarembo]

$$\langle W(C_{\text{circle}}) \rangle_{\lambda \gg 1} = \text{const} \frac{\lambda^3}{(\ln \lambda)^{3/2}}$$

that is equivalent to the semiclassical string in AdS_5

$$\langle W(C_{\text{circle}}) \rangle_{\lambda \gg 1} = KT^{-3/2} e^{2\pi T}$$

considering the string tension

$$T = \frac{3}{2\pi} \ln \lambda$$

AGT correspondence: The partition function of conformal $\mathcal{N} = 2$ theories on S^4 with gauge group $SU(N)$ is equivalent to a correlation function in A_{N-1} Toda CFT.

(A_1 Toda = Liouville)

[Alday, Gaiotto, Tachikawa]


AGT correspondence: The partition function of conformal $\mathcal{N} = 2$ theories on S^4 with gauge group $SU(N)$ is equivalent to a correlation function in A_{N-1} Toda CFT.

(A_1 Toda = Liouville)

[Alday, Gaiotto, Tachikawa]

Basic example: $SU(2)$ SCYM

$$Z_{SCYM} = \langle V_{\beta_1} V_{\beta_2} V_{\beta_3} V_{\beta_4} \rangle_{\text{Liouville}}$$



$$\int da^{(1)} Z_{\text{cl}} Z_{1\text{-loop}} Z_{\text{inst}} = \int d\alpha^{(1)} \langle \beta_1 | V_{\beta_2} | \alpha^{(1)} \rangle \langle \alpha^{(1)} | V_{\beta_3} | \beta_4 \rangle \mathcal{F}_{\alpha, \beta}(z) \bar{\mathcal{F}}_{\alpha, \beta}(\bar{z}) |z|^{2(\Delta_{\alpha} - \Delta_{\beta_3} - \Delta_{\beta_4})}$$


AGT correspondence: The partition function of conformal $\mathcal{N} = 2$ theories on S^4 with gauge group $SU(N)$ is equivalent to a correlation function in A_{N-1} Toda CFT.

(A_1 Toda = Liouville)

[Alday, Gaiotto, Tachikawa]

Basic example: $SU(2)$ SCYM

$$Z_{SCYM} = \langle V_{\beta_1} V_{\beta_2} V_{\beta_3} V_{\beta_4} \rangle_{\text{Liouville}}$$



$$\int da^{(1)} Z_{\text{cl}} Z_{1\text{-loop}} Z_{\text{inst}} = \int d\alpha^{(1)} \langle \beta_1 | V_{\beta_2} | \alpha^{(1)} \rangle \langle \alpha^{(1)} | V_{\beta_3} | \beta_4 \rangle \mathcal{F}_{\alpha, \beta}(z) \bar{\mathcal{F}}_{\alpha, \beta}(\bar{z}) |z|^{2(\Delta_{\alpha} - \Delta_{\beta_3} - \Delta_{\beta_4})}$$

- Different **S-duality frames** of the gauge theory correspond to the different **decomposition of the 2d CFT correlator**
- modular invariance of 2d CFT implies S-duality invariance of the 4d partition function

- Wilson, 't Hooft and dyonic line operators are realized as loop operators in Liouville/Toda CFT (Verlinde operators) [DGOT][AGGTT][Passerini][Gomis,LeFloch]
- modular transformations of loop operators in Toda CFT describe the action of S-duality on line operators in gauge theory
- for instance, it follows that $\langle \text{Wilson} \rangle_{\text{Theory}} = \langle \text{'t Hooft} \rangle_{S(\text{Theory})}$

- **Wilson, 't Hooft and dyonic line operators** are realized as loop operators in Liouville/Toda CFT (Verlinde operators) [DGOT][AGGTT][Passerini][Gomis,LeFloch]
- **modular transformations** of loop operators in Toda CFT describe the action of **S-duality** on line operators in gauge theory
- for instance, it follows that $\langle \text{Wilson} \rangle_{\text{Theory}} = \langle \text{'t Hooft} \rangle_{S(\text{Theory})}$

3d SUSY gauge theories have many interesting dualities: Mirror symmetry, AdS/CFT ...

- **partition functions** can be computed using **supersymmetric localizations** and many non-perturbative dualities can be tested explicitly [Kapustin, Willett, Yaakov]
- ABJM theory has the number of D.O.F. of M2 branes [Drukker, Mariño, Putrov]

In 3d, two main types of line operators: **Wilson loops and Vortex loops**

- **Vortex loop**: disorder line operator, defined by the path integral in the presence of a prescribed singularity along the defect line:
 - fixed gauge or flavor holonomy along infinitesimal curves linking the loop
 - singular matter fields

In 3d, two main types of line operators: **Wilson loops and Vortex loops**

- **Vortex loop**: disorder line operator, defined by the path integral in the presence of a prescribed singularity along the defect line:
 - fixed gauge or flavor holonomy along infinitesimal curves linking the loop
 - singular matter fields

the **VEV** of these operators can be computed using **supersymmetric localization**
[Drukker, Passerini, Okuda][Kapustin, Willett, Yaakov]

In 3d, two main types of line operators: **Wilson loops and Vortex loops**

- **Vortex loop**: disorder line operator, defined by the path integral in the presence of a prescribed singularity along the defect line:
 - fixed gauge or flavor holonomy along infinitesimal curves linking the loop
 - singular matter fields

the **VEV** of these operators can be computed using **supersymmetric localization**
[Drukker, Passerini, Okuda][Kapustin, Willett, Yaakov]

Interesting to study the role of line operators in dualities:

For instance, under **Abelian mirror symmetry**: **Wilson loops** \leftrightarrow **flavor vortex loops**

Conclusions

- supersymmetry provides a simplified framework to study Quantum Field Theory
- many novel tools for the weak and strong coupling dynamics of quantum fields (AdS/CFT, Pestun Localization, AGT, Integrability, ...)
- interconnections between gauge theory, string theory, two dimensional CFT and matrix models are useful to study the strong coupling dynamics of gauge theory and more formal aspects of string theory